

What is claimed is:

1. A vaccinating agent for use in promoting an effective immune response, in a mammalian host, against an infectious pathogen from the genus *Mycobacterium*, said vaccinating agent comprising:
 - 5 at least a portion of at least one majorly abundant extracellular product selected from the group consisting of *M. tuberculosis* 110 KD protein, 80 KD protein, 71 KD protein, 58 KD protein, 45 KD protein, 32A KD protein, 32B KD protein, 30 KD protein, 24 KD protein, 10 23.5 KD protein, 23 KD protein, 16 KD protein, 14 KD protein, 12 KD protein and respective analogs, homologs, and subunits thereof; and
 - an adjuvant selected from the group consisting of IL-12 and MF 59.
2. The vaccinating agent of claim 1 wherein said at least one majorly abundant extracellular product is *M. tuberculosis* 32A KD protein.
3. The vaccinating agent of claim 1 wherein said at least one majorly abundant extracellular product is a mixture of *M. tuberculosis* 32A KD protein, 30 KD protein, and 16 KD protein.
4. The vaccinating agent of claim 1 wherein said adjuvant is IL-12.
5. The vaccinating agent of claim 1 wherein said adjuvant is a mixture of IL-12 and MF 59.
6. A method for immunizing a mammalian host against an infectious pathogen of the genus *Mycobacterium*, said method comprising the steps of:
 - providing a vaccinating agent comprising at least a
 - 5 portion of at least one majorly abundant extracellular

267270-EE598/80

gird
up

product selected from the group consisting of *M. tuberculosis* 110 KD protein, 80 KD protein, 71 KD protein, 58 KD protein, 45 KD protein, 32A KD protein, 32B KD protein, 30 KD protein, 24 KD protein, 23.5 KD protein, 23 KD protein,
 10 16 KD protein, 14 KD protein, 12 KD protein and respective analogs, homologs, and subunits thereof, and an adjuvant selected from the group consisting of IL-12 and MF 59; and
 introducing said vaccinating agent into said
 mammalian host to induce an effective immune response to
 15 subsequent infection by said infectious pathogen.

7. The method of Claim 6 wherein said at least one majorly abundant extracellular product is *M. tuberculosis* 32A KD protein.

8. The method of claim 6 wherein said at least one majorly abundant extracellular product is a mixture of
M. tuberculosis 32A KD protein, 30 KD protein, and 16 KD
 protein.

9. The method of claim 6 wherein said adjuvant is IL-12.

10. The method of claim 6 wherein said adjuvant is a mixture of IL-12 and MF 59.

11. A vaccinating agent for use in promoting an effective immune response, in a mammalian host, against an infectious pathogen from the genus *Mycobacterium*, said vaccinating agent comprising:

at least one immunodominant epitope of at least one majorly abundant extracellular product selected from the group consisting of *M. tuberculosis* 110 KD protein, 80 KD protein, 71 KD protein, 58 KD protein, 45 KD protein, 32A KD protein, 32B KD protein, 30 KD protein, 24 KD protein, 23.5 KD protein, 23 KD protein, 16 KD protein,

14 KD protein, 12 KD protein, and respective analogs, homologs, and subunits thereof.

12. The vaccinating agent of claim 11 wherein said at least one majorly abundant extracellular product is *M. tuberculosis* 32A KD protein.

13. The vaccinating agent of claim 12 wherein said at least one immunodominant epitope is selected from the group consisting of *M. tuberculosis* 32A KD protein subunits having the amino acid sequences

	<u>Peptide Sequence</u>	<u>Seq. ID No.</u>
5	G L R A Q D D F S G W D I N T	104
	W D I N T P A F E W Y D Q S G	106
	P A F E W Y D Q S G L S V V M	107
	P V G G Q S S F Y S D W Y Q P	110
10	G C Q T Y K W E T F L T S E L	114
	K W E T F L T S E L P G W L Q	115
	A N R H V K P T G S A V V G L	118
	A V V G L S M A A S S A L T L	120
	S A L T L A I Y H P Q Q F V Y	122
15	A I Y H P Q Q F V Y A G A M S	123
	Q Q F V Y A G A M S G L L D P	124
	G L L D P S Q A M G P T L I G	126
	S Q A M G P T L I G L A M G D	127
	N D P L L N V G K L I A N N T	134
20	N V G K L I A N N T R V W V Y	135
	I A N N T R V W V Y C G N G K	136
	C G N G K P S D L G G N N L P	138

and respective analogs, homologs, and subunits thereof including single or multiple amino acid substitutions, deletions, insertions, and inversions.

2024-01-12 11:08am

14. An immunodiagnostic agent for use in promoting a detectable immune response in a mammalian host identifying an infectious pathogen from the genus *Mycobacterium*, said immunodiagnostic agent comprising:

5 at least one immunodominant epitope of at least one majorly abundant extracellular product selected from the group consisting of *M. tuberculosis* 110 KD protein, 80 KD protein, 71 KD protein, 58 KD protein, 45 KD protein, 32A KD protein, 32B KD protein, 30 KD protein, 24 KD
10 protein, 23.5 KD protein, 23 KD protein, 16 KD protein, 14 KD protein, 12 KD protein and respective analogs, homologs, and subunits thereof.

15. The immunodiagnostic agent of claim 14 wherein said at least one majorly abundant extracellular product is *M. tuberculosis* 32A KD protein.

16. The immunodiagnostic agent of claim 15 wherein said at least one immunodominant epitope is selected from the group consisting of *M. tuberculosis* 32A KD protein subunits having the amino acid sequences

	<u>Peptide Sequence</u>	<u>Seq. ID No.</u>
5	G L R A Q D D F S G W D I N T	104
	W D I N T P A F E W Y D Q S G	106
	P A F E W Y D Q S G L S V V M	107
	P V G G Q S S F Y S D W Y Q P	110
10	G C Q T Y K W E T F L T S E L	114
	K W E T F L T S E L P G W L Q	115
	A N R H V K P T G S A V V G L	118
	A V V G L S M A A S S A L T L	120
	S A L T L A I Y H P Q Q F V Y	122
15	A I Y H P Q Q F V Y A G A M S	123
	Q Q F V Y A G A M S G L L D P	124
	G L L D P S Q A M G P T L I G	126

087653-01497

	<u>Peptide Sequence</u>	<u>Seq. ID No.</u>
	S Q A M G P T L I G L A M G D	127
	N D P L L N V G K L I A N N T	134
20	N V G K L I A N N T R V W V Y	135
	I A N N T R V W V Y C G N G K	136
	C G N G K P S D L G G N N L P	138

and respective analogs, homologs, and subunits thereof including single or multiple amino acid substitutions, deletions, insertions, and inversions.

17. A method of immunizing a mammalian host against an infectious pathogen of the genus *Mycobacterium*, said method comprising the steps of:

providing at least one immunodominant epitope of at least one majorly abundant extracellular product selected from the group consisting of *M. tuberculosis* 110 KD protein, 80 KD protein, 71 KD protein, 58 KD protein, 45 KD protein, 32A KD protein, 32B KD protein, 30 KD protein, 24 KD protein, 23.5 KD protein, 23 KD protein, 16 KD protein, 14 KD protein, 12 KD protein and respective analogs, homologs, and subunits thereof; and

introducing said at least one immunodominant epitope to said mammalian host to induce an effective immune response to subsequent infection by said infectious pathogen.

18. The method of claim 17 wherein said at least one majorly abundant extracellular product is *M. tuberculosis* 32A KD protein.

19. The method of claim 18 wherein said at least one immunodominant epitope is selected from the group consisting of *M. tuberculosis* 32A KD protein subunits having the amino acid sequences

08786533-01419

	<u>Peptide Sequence</u>	<u>Seq. ID No.</u>
5	GLRAQDDFSGW DINT	104
	WDINTPAFEWYDQSG	106
	PAFEWYDQSGLSVVM	107
	PVGGQSSSFYSDWYQP	110
10	GCQTYKWETF LTSEL	114
	KWETF LTSELPGWLQ	115
	ANRHVKPTGSAVVGL	118
	AVVGLSMAASSALT L	120
	SALT LAIYHPQQFVY	122
15	A IYHPQQFVYAGAMS	123
	QQFVYAGAMSGLLDP	124
	GLLDPSQAMGPTLIG	126
	SQAMGPTLIGLAMGD	127
	NDPLLN VGKLIANNNT	134
20	NVGKLIANNTRVWVY	135
	IANNTTRVWVYCGNGK	136
	CGNGKPSDLGGNNLP	138

and respective analogs, homologs, and subunits thereof including single or multiple amino acid substitutions, deletions, insertions, and inversions.

20. A method for detecting the presence of an immune response in a mammal against a pathogen of the genus *Mycobacterium*, said method comprising the steps of:

- providing at least one immunodominant epitope of
- 5 at least one majorly abundant extracellular product selected from the group consisting of *M. tuberculosis* 110 KD protein, 80 KD protein, 71 KD protein, 58 KD protein, 45 KD protein, 32A KD protein, 32B KD protein, 30 KD protein, 24 KD protein, 23.5 KD protein, 23 KD protein, 16
- 10 KD protein, 14 KD protein, 12 KD protein and respective analogs, homologs, and subunits thereof;

administering said at least one immunodominant epitope to said mammal; and
measuring the resultant immune response.

21. The method of claim 20 wherein said at least one majorly abundant extracellular product is *M. tuberculosis* 32A KD protein.

22. The method of claim 21 wherein said at least one immunodominant epitope is selected from the group consisting of *M. tuberculosis* 32A KD protein subunits having the amino acid sequences

	<u>Peptide Sequence</u>	<u>Seq. ID No.</u>
5	G L R A Q D D F S G W D I N T	104
	W D I N T P A F E W Y D Q S G	106
	P A F E W Y D Q S G L S V V M	107
	P V G G Q S S F Y S D W Y Q P	110
10	G C Q T Y K W E T F L T S E L	114
	K W E T F L T S E L P G W L Q	115
	A N R H V K P T G S A V V G L	118
	A V V G L S M A A S S A L T L	120
	S A L T L A I Y H P Q Q F V Y	122
15	A I Y H P Q Q F V Y A G A M S	123
	Q Q F V Y A G A M S G L L D P	124
	G L L D P S Q A M G P T L I G	126
	S Q A M G P T L I G L A M G D	127
	N D P L L N V G K L I A N N T	134
20	N V G K L I A N N T R V W V Y	135
	I A N N T R V W V Y C G N G K	136
	C G N G K P S D L G G N N L P	138

and respective analogs, homologs, and subunits thereof including single or multiple amino acid substitutions,
25 deletions, insertions, and inversions.

08786533-012197

23. A process for producing a majorly abundant extracellular product selected from the group consisting of *M. tuberculosis* 110 KD protein, 80 KD protein, 71 KD protein, 58 KD protein, 45 KD protein, 32A KD protein, 32B
 5 KD protein, 30 KD protein, 24 KD protein, 23.5 KD protein, 23 KD protein, 16 KD protein, 14 KD protein, 12 KD protein and respective analogs, homologs, and subunits thereof, said process comprising the steps of:

transforming a host cell with a vector to form
 10 a transformed cell, said vector comprising a nucleic acid molecule encoding one of said majorly abundant extracellular products; and

culturing said transformed cell to thereby produce said majorly abundant extracellular product.

24. The process of claim 23 wherein said nucleic acid molecule encodes for the 32A KD *M. tuberculosis* protein.

25. The process of claim 24 which includes the additional step of recovering said majorly abundant extracellular product that is produced by culturing of said transformed cell.

26. The process of claim 24 wherein said vector comprises pSMT3. LAB designation

27. The process of claim 24 wherein said host cell is *M. smegmatis* or *M. vaccae*.

28. The process of claim 24 wherein said transformed cell is cultured at a temperature of 28°C.

08786533-012197